

Mortality

The Effects Of Harmful Alcohol Consumption

13.3 million killed

accounting for about

5.9% of all deaths1

Number 1 risk factor for premature death in men aged 15 to 59 years²

70% of net mortality


caused by harmful alcohol consumption and alcohol use disorders (AUDs) in Europe³

Alcohol is a risk factor to over 60 types of diseases⁵



5.1%

of the global burden of disease⁴





Up to 12% of European children live in families adversely affected by alcohol8


Costs due to alcohol in the EU, 2010 (€ billion)10



Costs

€155.8 billion

Costs of excessive alcohol consumption, 60% of which are attributable to the harmful use alcohol and AUDs⁹




1/4
of workplace accidents
linked to alcohol¹¹

Healthcare interventions for harmful alcohol consumption

Low risk Hazardous Harmful drinking drinking drinking (>20 grams per day (>20 grams per day (<20 grams per Severe disorder day - WHO AUDIT WHO AUDIT score WHO AUDIT score (>20 grams per day – WHO AUDIT score 40-20) score 7-0) 15-8) 19-16) Monitor your Brief Treatment in Specialized treatment drinkina intervention primary care Higher consumption of alcohol


Alcohol can hijack your brain: the harmful alcohol use cycle






The Effects Of Harmful Alcohol Consumption

Alcohol is toxic to most organs: the harmful effects of alcohol on the human body^{13,14}





*Source: https://www.ima.org.il/FilesUpload/IMAJ/0/55/27730.pdf

- WHO, 2014, Status Report on Alcohol, http://www.who.int/iris/bitstream/10665/112736/1/9789240692763 enq.pdf?ua=1
- European Commission, 2011, Alcohol, Work and Productivity: Scientific Opinion of the Science Group of the European Alcohol and Health Forum, http://ec.europa.eu/health/alcohol/docs/science_02_en.pdf
- Rehm J and Shield KD, 2012, Interventions for alcohol dependence in Europe: a missed opportunity to improve public health, based on the book "Alcohol consumption, alcohol dependence and attributable burden of disease in Europe: potential gains from effective interventions for alcohol dependence
- WHO, 2014, op. cit.
- WHO, 2010, Status Report on NCDs, http://www.who.int/iris/bitstream/10665/112736/1/9789240692763_eng.pdf?ua=1
- OECD, 2013, Harmful Use Of Alcohol: Trends And Policy Conclusions, Directorate For Employment, Labour And Social Affairs' Health Committee
- NICE National Clinical Practice Guidelines 115, 2011, Alcohol-use disorders. Diagnosis, assessment and management of harmful drinking and alcohol dependence. Page 34
- Anderson & Baumberg, 2006, A report for the European Commission, Institute of Alcohol Studies; Alcohol in Europe. A Public Health Perspective. Page 203
- Rehm J and Shield KD, 2012, op. cit.
- Rehm J et al., 2012, Modeling the impact of alcohol dependence on mortality burden and the effect of available treatment interventions in the European Union, Schuckit Ch. 98. In: Davis et al. Neuropsychopharmacology: The fifth Generation of Progress
- 2002. Page 1399
 European Workplace and Alcohol project (EWA), 2013, Factsheet http://www.eurocare.org/content/download/15763/87219/file/EWA%20Factsheet%20Final.pdf
- Rehm J, 2014, Centre for Addiction and Mental Health, Toronto, Canada. Expert Roundtable Meeting: Screening and intervention for harmful alcohol use as a tool to improve the management of hypertension in primary care, 06 May 2014 Barcelona.
- 13. http://drink-less.com/global/whats-at-stake/effects-on-the-body-men
- http://drink-less.com/global/whats-at-stake/effects-on-the-body-women Anderson P and Baumberg B, 2006, A report for the European Commission, Institute of Alcohol Studies; Alcohol in Europe, A Public Health Perspective, Page 142
- Rehm & Roerecke, 2013, Alc Alc
- 17. Rehm & Roerecke, 2013, Alc Alc